


## Behavioral Emergencies - Special Considerations in the Pregnant Patient



**Across**

- 3. \_\_\_\_\_ also appears to be more likely to occur during pregnancy compared with the postpartum period.
- 5. A 2013 meta-analysis of 19 studies reported no association between \_\_\_\_\_ exposure and congenital malformations, as well as major malformations.
- 6. The severity of agitations may warrant appropriate clinical use of \_\_\_\_\_ after the risks and benefits of short-term administration have been weighed
- 7. Women in the second and third trimesters of pregnancy are vulnerable to the development of \_\_\_\_\_ if mechanical restraints are used.
- 9. Although methadone remains the current standard of care, \_\_\_\_\_ is increasingly being preferred by many practitioners, given limited access to methadone maintenance treatment programs, similar efficacy, and other relatively favorable outcomes described in this article.

- 10. Maternal opioid withdrawal can lead to the development of an \_\_\_\_\_ abstinence syndrome, which could be potentially fatal for the fetus, and can have potential longterm effects on the development of the fetal brain.

**Down**

- 1. \_\_\_\_\_ can be of benefit in managing mild alcohol withdrawal syndrome, and may be a consideration, given its relative safety in pregnancy.
- 2. The risk of neural tube defects with \_\_\_\_\_ is significantly high but less than that of valproate.
- 4. The \_\_\_\_\_ properties of alcohol have been recognized since the 1960s.
- 8. A prior history of \_\_\_\_\_, perinatal or otherwise, is the biggest factor predictive of future risk

**Word Bank**

- |                |                    |              |
|----------------|--------------------|--------------|
| Suicidality    | benzodiazepines    | Depression   |
| antidepressant | teratogenic        | intrauterine |
| buprenorphine  | vena cava syndrome | Gabapentin   |
| carbamazepine  |                    |              |