


Name: \_\_\_\_\_

Date: \_\_\_\_\_

# Brain Injury



## Across

3. One of the best ways to protect our brain is to wear a \_\_\_\_\_.

6. A person who sustains a brain injury and loses consciousness for longer than 30 minutes and has confusion for longer than 24 hours has a \_\_\_\_\_ brain injury.

7. To keep a Brain Injured patient from becoming overstimulated and agitated, we should keep the patient's room \_\_\_\_\_.

8. The most common cause of traumatic brain Injuries is \_\_\_\_\_.

11. A mild brain injury is also often called a \_\_\_\_\_.

12. An Injury to the brain that is not congenital, hereditary, degenerative, or induced by birth trauma is an \_\_\_\_\_ Brain Injury.

## Down

1. The age group most likely to sustain a traumatic brain injury is age \_\_\_\_\_ and older.

2. \_\_\_\_\_ are more likely to sustain a Brain Injury.

4. A person who sustains a brain injury that does not lose consciousness or loses consciousness for less than 30 minutes and has confusion for less than 24 hours has a \_\_\_\_\_ brain injury.

5. An Injury to the brain from an external force is a \_\_\_\_\_ Brain Injury.

9. If a brain injured patient becomes agitated and staff needs assistance, they should call a code \_\_\_\_\_.

10. When interacting with a brain injury patient, you should speak \_\_\_\_\_.