


Cardiac



Across

2. A medically treated 4.4 cm TAA can have _____ follow-up imaging.

8. T wave inversion in lead v5 is an _____ EKG.

11. PVD-3.1 would be used to evaluate a new _____ bruit.

13. Multigated Acquisition scan

15. A patient who meets all of the criteria for an imaging stress test can also have a cardiac PET if he/she is _____ obese.

16. Blood test used to diagnose and track heart failure.

18. A marker for ischemic injury; primarily cardiac.

19. Test used to determine blood sugar control for patients with diabetes.

Down

1. Shortness of breath when upright or seated; the opposite of orthopnea.

3. Pre-test probability of CAD is determined by age, _____, and symptoms.

4. _____ vena cava (SVC)

5. Initial imaging study to evaluate for deep venous thrombosis (DVT)

6. Painful cramping sensation of the legs with walking or severe leg fatigue

7. A 93306 is not indicated in the eval of _____ congenital heart disease.

9. _____ chest discomfort is a sign of typical angina.

10. A cardiac CT, CTV, MRV, or MRI can be used to eval pulmonary _____.

12. _____ syncope is NOT a reason to approve a left heart catheterization.

14. Loss of consciousness

17. Greater than 90% pre-test probability of CAD is considered "_____" probability.