


# Ch 61 & 62



## Across

3. When low omphalocele is observed, consider bladder or cloacal \_\_\_\_\_.
5. Exposed heart presents outside chest wall through cleft sternum is known as what?
9. Intestines return to abdominal cavity by \_\_\_\_\_ week of gestation.
11. In 6th week, process called \_\_\_\_\_ helps embryo transform itself into cylindrical shape.
12. The connection of the \_\_\_\_\_ and body stalk will form the umbilical cord
15. 2 of 3 layers by end of 5th week of development,
17. High or \_\_\_\_\_ omphalocele usually primary finding

18. Most commonly occurs from prolonged \_\_\_\_\_ or secondary to small thoracic cavity as result of structural or chromosomal abnormality
19. 2 of 3 most common abdominal wall defects
20. 1 of 3 layers by end of 5th week of development,
- ## Down
1. in this anomaly, \_\_\_\_\_ organs form complex, bizarre-appearing mass entangled with membranes
2. A \_\_\_\_\_ is the most common lung cyst detected
4. When \_\_\_\_\_ is found, consider limb-body wall complex
6. 1 of 3 most common abdominal wall defects

7. Accumulation of fluid within pleural cavity is called what?
8. 3 of 3 most common abdominal wall defects
10. 3 of 3 layers by end of 5th week of development,
13. Dramatic pulsations of anterior chest wall occur from heart beating against chest without presence of the \_\_\_\_\_.
14. \_\_\_\_\_ grows faster than abdomen.
16. \_\_\_\_\_ development single most determinate of viability.