


Colon Cancer



Across

- 5. This blood test is the specific tumor marker for colorectal cancer
- 7. The main test use to diagnose colorectal cancer
- 10. What type of risk factor are alcohol intake and smoking considered?
- 12. How many people are estimated to have died from colon cancer in Canada in 2017?
- 13. The common first line chemotherapy treatment for stage 4 colorectal cancer is FOLFIRI, which is given ever ___ weeks.
- 14. Individuals diagnosed with stage 2 cancer often receive adjuvant chemotherapy to reduce risk of recurrence that includes oxaliplatin, leucovorin, and 5-fluorouracil. What is the common acronym for this treatment?

- 15. Iv therapy for the treatment of cancer is called?
 - 16. If cancer has spread to other areas of the body, it it called _____ bowel cancer
 - 17. Which of the 5 common chemotherapy medications used to treat colon cancer are missing? Irinotecan, 5-fluorouracil, capecitabine, raltitrexed
- Down**
- 1. This is the stage of cancer if there are cancer cells noted in the lymph nodes near the colon (Roman numerals)
 - 2. _____ is the most common type of colorectal cancer
 - 3. This at home test can detect early colorectal cancer
 - 4. The test often used to determine if there is metastatic disease in the lymph nodes or other abdominal organs.

- 6. This bowel disease causing sores and bleeding, can increase risk of bowel cancer
- 8. _____ chemotherapy is given to a patient after the tumor is surgically removed to prevent recurrence .
- 9. This test take a sample of tissue within the colon to test for presence of cancer cells
- 11. How many cases of colon cancer were diagnosed in Canada in 2017?