


Name: \_\_\_\_\_

Date: \_\_\_\_\_

# Dentistry Crossword



**Across**

- 3. AAHA guideline recommends not using \_\_\_\_\_ which can cause myalgia, neuralgia, and/or trauma to the temporomandibular joint.
- 6. A feline permanent dental formula shows an absence of the \_\_\_\_\_ maxillary premolar.
- 11. \_\_\_\_\_ is the correct term referring to the surface of the tooth facing the vestibule or lips. Buccal or labial are acceptable alternatives.
- 12. Lidocaine and \_\_\_\_\_ are the common regional anesthetic agents used for dental procedures.
- 13. Prevalence of feline odontoclastic resorptive oral lesions increase with \_\_\_\_\_.
- 14. Gingival enlargement is known to be associated with \_\_\_\_\_ channel blockers.
- 17. \_\_\_\_\_ exposure exists when a periodontal probe extends under the crown of a multirooted tooth, through and through from one side of the furcation out the other.

- 18. Stage \_\_\_\_\_ periodontal disease is classified by less than 25% of attachment loss.
  - 19. \_\_\_\_\_ is the term used to describe tooth wear caused by contact of a tooth with a non-dental object.
  - 20. Ingestion of \_\_\_\_\_ during tooth development can cause generalized intrinsic staining of teeth.
- Down**
- 1. A \_\_\_\_\_ tooth is the presence of an extra tooth.
  - 2. Ideally, irreversible pulpitis and secondary pulp necrosis could be definitively diagnosed based on \_\_\_\_\_.
  - 4. \_\_\_\_\_ is the term used to describe a disturbance in tooth development that causes the crown or root to be abruptly bent or crooked.
  - 5. The \_\_\_\_\_ surface of of the canines, premolars, and molars is the surface that points towards the first incisor.

- 7. \_\_\_\_\_ are an increased probing depth of a tooth even in the absence of attachment loss.
- 8. A \_\_\_\_\_ cyst is formed around the crown of a partially erupted or unerupted tooth.
- 9. Class \_\_\_\_\_ malocclusion is an abnormal rostralcaudal relationship between the dental arches in which the mandibular arch occludes rostral to its normal position relative to the maxillary arch.
- 10. \_\_\_\_\_ itself is not desirable as a sole agent for local block due to its limited effect post administration (1-2 hours).
- 15. The teeth that are most commonly affected with feline odontoclastic resorptive lesions are \_\_\_\_\_ third premolars.
- 16. The right maxillary deciduous arcade begins with the number \_\_\_\_\_.