


# Diabetes hypoawareness



## Across

2. You can only give this drug once in 24 hours intramuscularly (8)  
 6. C<sub>2</sub>H<sub>5</sub>OH is a simple example of the chemical that may induce hypoglycaemia events (7)  
 11. Two tubes of this is given orally for treatment (8)  
 13. 5 of these recommended sweets can be used as treatment (5,6)  
 14. This class of drug can cause hypoglycaemic events (14)  
 15. Total lack of insulin results in this condition (4,3,8)  
 18. Operation - usually performed for pancreatic cancer (8)  
 19. A word for when a person has no signs or symptoms of a hypoglycaemic episode (7)

20. Insulin is called as this (7)  
 21. Impairment of this organ may result in hypoglycaemia events (7)  
 22. Reducing this class of drug can result in hypoglycaemia events (8)  
 23. You might experience this during a hypoglycaemic episode (8)

## Down

1. Simple sugar (7)  
 3. Medical term for blood glucose below four millimoles per litre (13)  
 4. Complex sugar group given after a hypoglycaemic event (13)  
 5. Recommended time between the next blood glucose test (7,7)  
 7. Accumulation of extra fat at injection sites that impairs insulin absorption (15)

8. Insulin is produced, stored and released here (4,5)  
 9. Refer to the diabetes team using this (6)  
 10. Drink given initially to a person with a low blood glucose (6,5)  
 12. Safe level of blood glucose for a person to drive if on insulin injections (4)  
 16. An app with our diabetes UHS guidelines (10)  
 17. The organ insulin is associated with (8)