


Gastroenteritis Complications



Across

- 2.** the toxin produced by a subclass of STEC that can cause HUS
3. the disease most commonly associated with Toxic Megacolon
6. the most common complication seen in patients with diarrhea
8. the most common infectious cause of Toxic Megacolon
10. The percentage of patients that report a viral infection before diagnosis of Intussusception
11. a class of E.Coli that cause severe diarrhea

- 12.** the vaccine that was associated with a 22 fold increase in Intussusception among children thirty, the percentage of patients
13. bacteria associated with higher risk for systemic infection (pneumonia, meningitis) and sepsis
14. a complication of diarrhea that can cause rapid weight loss
15. Must be present in a triad of symptoms, with nonimmune mediated hemolytic anemia and acute kidney injury to diagnose HUS
17. telescoping of the intestine into itself that can occur at multiple points throughout the GI tract

Down

- 1.** a syndrome seen mostly in children infected with STEC, that can cause acute kidney injury
4. a disorder caused by rapid loss of bicarb into the lumen of the GI tract
5. STEC infections commonly present with abdominal pain, vomiting, bloody diarrhea, but no _____
7. total or segmental non-obstructive colonic dilation plus systemic toxicity
9. the electrolyte most susceptible to large changes in serum concentration with diarrhea and vomiting
16. a complication of diarrhea seen in adults and associated with Salmonella infection