


HIV and STD



Across

- 2. sometimes called the "Clap" or the "Drip" _____
- 5. Condoms should not be used with lubricants that contain this substance _____
- 6. The most common STI in the United States, it can cause genital warts and cervical cancer _____
- 9. STI and STD's are _____ that are passed from one person to another through sexual contact.
- 11. It is the virus that causes AIDS _____.
- 12. This is the test that detects presence of mutations associated with ART drug resistance _____

- 14. A viral load of 50 copies/ml or less is considered _____
- 17. Most effective form of birth control and disease prevention _____.

Down

- 1. The virus that causes sores or blisters that appear on the genitals of both male and females is known as _____
- 3. Term for taking antiretrovirals (ART) medications as soon as possible after exposure to HIV., so that exposure will not result in HIV is known as Post _____ Prophylaxis
- 4. Another name for STI _____

- 7. Three ways mother to child transmission can occur is in utero, birth and _____milk.
- 8. What is the acronym for Acquired Immune Deficiency? _____
- 10. CD4+ Cells (Tcells) are the _____ cells/soldiers of the immune system
- 13. If a person experiences burning or pain with urination it could be an _____
- 15. Condoms that are made of polyurethane and _____is effective in the prevention of HIV.
- 16. Sharing needles for tattooing or ear piercing is high _____.