


Heat Illness Awareness



Across

- 3.** High Humidity, Lack Of Air Movement, Air Temperature, Radiant Heat Temperature, & Ground Temperature are ? Risk Factors
- 8.** Consumption Of ? Essential Prior To Conducting New Tasks, Operating Equipment, & Donning Tyvek/Tychem Suit
- 10.** Mandatory Every 2 Hours During High Heat Procedures
- 11.** Without Outside Intervention Heat Stroke Can Be ?

- 12.** Sweaty Skin, Fast Heart Beat, Cramps, Headache, Weakness, Dizziness, Nausea, & Vomiting Are Symptoms Of ?

Down

- 1.** Remember: If you feel thirsty you are already ?
- 2.** 80% Of The Reported Cases of Heat Illness Occurred Within The First ? Days
- 4.** Red, Hot Dry Skin, Confusion, Convulsion, High Temperature, Fainting

- 5.** Age, Medication, Weight Physical Fitness, Caffeine/Alcohol Consumption, & Overexertion Are ? Risk Factors
- 6.** Exceeding 80 Degrees And At Least 10 Degrees Higher Than Preceding Five Days
- 7.** Takes 5-14 Days With Consistent High Heat
- 9.** 40% Of The Reported Cases of Heat Illness Occurred On the Employee's ? Day
- 13.** Made Available At 80 Degrees Or Higher