


Name: \_\_\_\_\_ Date: \_\_\_\_\_

### Ineffective Peripheral Tissue Perfusion Secondary to Atherosclerosis and Right Leg Claudication



#### Across

- 4. Decreased \_\_\_\_\_ is a sign of ineffective tissue perfusion.
- 5. Pain caused by too little blood flow, usually during exertion.
- 7. Should be encouraged soon after procedure to promote circulation.
- 8. An antiplatelet medication that should be continued after procedure.
- 9. Changes in \_\_\_\_\_ temperature, color, or sensation, or the presence of a new lesion on the lower extremities should be reported to PCP.

- 10. Diabetes, smoking, eating habits, level of exercise, low HDL, high LDL, and genetic predisposition.

#### Down

- 1. Plaque forms in the arterial walls causing thickening and hardening of the vessels.
- 2. Patient admitted with right leg claudication and premature vascular disease.
- 3. When the patient is discharged with \_\_\_\_\_ educate on checking for proper suction and knowing how to reinforce dressing.
- 6. Use of a \_\_\_\_\_ can aide in checking pedal pulses.

#### Word Bank

- |              |                 |            |                 |
|--------------|-----------------|------------|-----------------|
| Skin         | Chief complaint | Doppler    | Plavix          |
| Wound vac    | Comorbidities   | Ambulation | Atherosclerosis |
| Claudication | Pedal pulses    |            |                 |