


# MOS



## Across

2. Another name for stitches  
 3. These forceps are used for extraction of an upper wisdom tooth  
 7. This medicament (gel) is used prior to using local anaesthetic  
 9. This type of needle is required for infiltration  
 10. Breech loading syringe  
 12. Name given when bleeding has stopped  
 13. Anaesthetic liquid used to numb the patients mouth  
 18. This type of bleeding happens 24 hours after an extraction

20. This type of bleeding happens 6-8 hours after extraction

21. This item is used to make an incision into soft tissues  
 22. Side loading syringe  
 23. Vasoconstrictor in local anaesthetic and ingredient found in haemostatic packs
- ## Down
1. How long should a patient wait before rinsing after a tooth extraction (--hours)  
 4. Roots are that left behind in a socket  
 5. A piece of tissue that is removed for investigation purposes

6. Written instructions provided after a tooth extraction

8. Name given to infection of a socket after tooth extraction  
 11. Used to bite on after an extraction  
 14. This type of needle is required for block nerve injections  
 15. Acronym used for tooth extraction on a dental record  
 16. Inflammation of the soft tissues surrounding the crown of a partially erupted tooth  
 17. This type of waste disposal is required for needles  
 19. This type of haemorrhage happens immediately after extraction