


Neonatal/Pediatric Respiratory Care



Across

2. Occurs when a placenta partially or totally covers mother's cervix
 6. A scaphoid abdomen is most often associated with
 8. APGAR Score 4-6
 9. Administered to stimulate surfactant production before birth
 11. P in APGAR
 19. Number of pregnancies resulting in a potentially viable infant
 20. Number of pregnancies
 22. Placenta partially or completely separates from inner wall of uterus before delivery
 24. R in APGAR

25. Treatment for jaundice

26. APGAR Score 0-3
 27. What is missing in the lungs that causes RDS

28. Normal 120-160

Down

1. What would an x-ray showing "egg shaped heart" indicate
 3. Normal 40-60
 4. PROM
 5. Blood vessel present in all babies while still in the womb, allows blood to bypass lungs
 7. Pharmacological treatment to encourage closure of PDA

10. Refers to a woman's first pregnancy

12. Reported at 1 and 5 minutes after birth

13. Abnormal opening in the dividing wall between upper filling chambers of the heart

14. What would a Dubowitz score of 34 indicate?

15. X-ray technique to confirm diagnosis of epiglottitis

16. Pharmacological agent used to keep the PDA open

17. 2nd A in APGAR

18. A in APGAR

21. G in APGAR

23. APGAR Score 7-10