


Perinatal Final



Across

2. painless dilation of cervix without contractions
4. Should not be used with other calcium channel blockers
5. Unilateral abd pain, fainting, dizziness and referred right should pain
12. Rupture before 37weeks
13. Medications to help stop labor. Delays 24-48hrs
15. contractions lasting longer than 2 mins
17. primipara cervical dilation per hour
21. New onset of HTN with protein uria and edema
22. Signs include depression or absence of reflexes, oliguria, confusion, respiratory depression, respiratory paralysis.
23. One or more lobes developed. PP hemorrhage if minor lobe remains in uterus
24. Bacterial infection found in the lower GI tract. Leading cause of neonatal sepsis and mortality
25. Vaginal bleeding and passage of grapelike clusters of tissue
28. Artificial rupture of membrane

29. Painless bleeding and 4 types Total, partial, marginal and low lying

30. 15x15 for 20 min period

31. Persistent vomiting past the first trimester that can result in deficiency and dehydration

33. Three contractions of good quality lasting 40 or more seconds in 10 min without evidence of late decelerations

34. Fetal surface of the placenta is exposed through a ring opening around umbilical cord

Down

1. Tool used to determine ectopic pregnancy, FHR, fetal life and amniotic fluid

3. Pain in abdomen, back and pelvis; menstrual cramps; vaginal bleeding; increased vaginal discharge; pelvic pressure; urinary frequency; diarrhea

6. Procedure to detect genetic, metabolic, and DNA abnormalities

7. ROA, LOA

8. Glycoprotein produced by the trophoblast and other fetal tissues. Absence between 20-34weeks strong predictor of non preterm birth

9. Threatened, Imminent, Incomplete, Complete and Missed are types of

10. abnormal labor patterns occur with power, passenger and passage

11. FHR acceitation, Breathing, Tone, Movement and Amniotic fluid

14. Mg Sulfate reversal agent

16. Marginal, central and complete

18. Frank, footling and complete

19. Newborn weight >90th percentile

20. >42weeks

26. 39-40.6 weeks

27. Administered at 28 weeks gestation and 72 hours after birth

32. <10th percentile growth curve