


# Peripheral Artery Disease



**Across**

- 2. Cardinal sign of PAD
- 3. Who you would consult for the toe ulcer
- 8. HTN medication used for this patient
- 9. 6Ps that involves decreased movement
- 13. Most common location of arterial ulcers
- 17. 6Ps that involves cold extremities
- 18. 6Ps that involves discomfort and hurting
- 19. Patient Age
- 20. 6Ps that involves white-ish color to the skin
- 21. Priority body system for PAD that involves circulation
- 22. 6Ps that involves no palpable pulse

- 25. Priority body system for PAD that involves skin
  - 26. Patient Name
  - 27. Providing patient with information about modifiable risk factors
  - 28. Education about how to stop smoking
  - 29. Diagnostic that compares blood pressure in arm to that in leg
- Down**
- 1. When the legs are elevated, blood can't get down to the lower extremities
  - 4. Education about movement and weight loss
  - 5. X-ray that can show blockage or narrowing of arteries
  - 6. Medication used to manage diabetes
  - 7. 6Ps that involves decreased sensation or tingling

- 10. Biggest risk factor of PAD
- 11. Medication that relieves symptoms of intermittent claudication
- 12. Risk factor of PAD that results in high blood sugar
- 14. Diet that involves limiting sodium and fat intake
- 15. Ultrasound that can show decreased blood flow
- 16. Elevated blood pressure
- 23. Cholesterol medication used for this patient
- 24. Pain medication used for this patient