


Pharmacology week 2 blank questions



Across

9. Intramuscular injections for infants should be given below the greater trochanter of the femur but within the upper _____ of the vastus lateralis muscle.

10. A tablet can be cut in half only if the manufacturer has prepackaged it as a(n) _____ tablet.

11. Antibiotics typically are prescribed initially in a(n) _____ dose to reach therapeutic blood levels as quickly as possible.

12. A tablet may be given a(n) _____ coating so that the medicine is absorbed in the small intestine.

13. At the dorsogluteal intramuscular site, a primary concern is to avoid injecting the medication into the _____ nerve.

14. A medication to open the airways is prescribed because the patient has _____.

15. A(n) _____ is a bottle of medication with a rubber stopper that contains enough medication for more than one injection.

Down

1. Repeated injections of minute amounts of allergens to desensitize a patient to allergic reactions is called _____.

2. _____ are administered to patients with peripheral edema to help absorb extra fluid from cells.

3. Subcutaneous injections are administered just under the dermal layers into fatty, or _____, tissue.

4. At the dorsogluteal intramuscular site, the injection is administered into the _____ muscle.

5. In an infant, the safest site for an intramuscular injection is the _____.

6. The _____ of a medication is its thickness and its ability to move easily from a vial to an injection unit.

7. _____ is an enlarged, hardened reaction produced by the Mantoux test.

8. If an IV cannula becomes dislodged from a blood vessel, _____ of tissues at the site can occur.