


Name: \_\_\_\_\_

Date: \_\_\_\_\_

# Phlebotomy



## Across

1. \_\_\_\_\_ is the destruction of red blood cells  
8. \_\_\_\_\_ may lead to inaccurate test results  
9. Capillary sticks are performed on a infants \_\_\_\_\_  
10. You do not have to wear \_\_\_\_\_ gloves for a phlebotomy stick.  
11. When performing blood glucose check, the first drip of blood is \_\_\_\_\_ away.

12. The \_\_\_\_\_ thumb finger has its own pulse and should not be used for a capillary stick  
14. Always dispose of needles in the \_\_\_\_\_ container  
15. The tourniquet can not stay on longer than \_\_\_\_\_ minute.

## Down

2. Always remove the \_\_\_\_\_ before the needle.  
3. \_\_\_\_\_ is separated from whole blood in a SST tube once centrifuge.

4. A butterfly is also known as a \_\_\_\_\_ set.  
5. the most common space for a venipuncture is the \_\_\_\_\_ region.  
6. If a capillary stick and not be done on a patient finger or heel, you can use their \_\_\_\_\_.  
7. A hemoglobin test can be used to check \_\_\_\_\_.  
13. The \_\_\_\_\_ of the needle should always be up when performing a venipuncture