

Phlebotomy

1. Blood culture tubes, no inversions, can be done to r/o infections
 2. Coagulation tubes, 4 inversions additive Sodium Citrate
 3. Often used in blood bank(cross match) immunology and seriology
 4. used for PT PTT has Heparin additive
 5. Has EDTA additive, 8 inversions, used for CBC, WBC, RBC
 6. This is used to preserve glucose in whole blood, also for special chemistry tests
 7. Red blood cells per microliter of blood
 8. Heel sticks performed on
 9. The 2 hr postprandial test is used for
 10. Prandial
 11. found in capillaries
 12. normal range for insulin testing
 13. hypoglycemia
 14. Normal hemoglobin for female
 15. normal hemoglobin for males
 16. Normal fasting glucose level
 17. normal hematocrit in males
 18. Lab testing conducted close to the site of patient care
 19. Good cholesterol
 20. Normal total cholesterol
- A. POCT
 - B. red top tube
 - C. diabetes mellitus
 - D. HDL
 - E. Yellow top tube
 - F. 13.5-17.5
 - G. fasting
 - H. Lavender top tube
 - I. 4.2-6.2 billion
 - J. blue top tube
 - K. low blood sugar
 - L. <17UmL
 - M. venous and arterial blood
 - N. 60-110
 - O. >200mg/dl
 - P. green top tube
 - Q. 12.0-15.0
 - R. infants less than 1 yr
 - S. gray top tube
 - T. 40-55%